

# UNIVERSITY OF BIRMINGHAM

University of Birmingham  
Research at Birmingham

## Postnatal Deletion of Podoplanin in Lymphatic Endothelium Results in Blood Filling of the Lymphatic System and Impairs Dendritic Cell Migration to Lymph Nodes

Bianchi, Roberta; Russo, Erica; Bachmann, Samia B; Proulx, Steven T; Sesartic, Marko; Smaadahl, Nora; Watson, Steve P; Buckley, Christopher D; Halin, Cornelia; Detmar, Michael

DOI:

[10.1161/ATVBAHA.116.308020](https://doi.org/10.1161/ATVBAHA.116.308020)

License:

None: All rights reserved

*Document Version*

Peer reviewed version

*Citation for published version (Harvard):*

Bianchi, R, Russo, E, Bachmann, SB, Proulx, ST, Sesartic, M, Smaadahl, N, Watson, SP, Buckley, CD, Halin, C & Detmar, M 2016, 'Postnatal Deletion of Podoplanin in Lymphatic Endothelium Results in Blood Filling of the Lymphatic System and Impairs Dendritic Cell Migration to Lymph Nodes', *Arteriosclerosis Thrombosis and Vascular Biology*, vol. 36, no. 12. <https://doi.org/10.1161/ATVBAHA.116.308020>

[Link to publication on Research at Birmingham portal](#)

### **Publisher Rights Statement:**

Eligibility for repository: Checked on 30/11/2016

### **General rights**

Unless a licence is specified above, all rights (including copyright and moral rights) in this document are retained by the authors and/or the copyright holders. The express permission of the copyright holder must be obtained for any use of this material other than for purposes permitted by law.

- Users may freely distribute the URL that is used to identify this publication.
- Users may download and/or print one copy of the publication from the University of Birmingham research portal for the purpose of private study or non-commercial research.
- User may use extracts from the document in line with the concept of 'fair dealing' under the Copyright, Designs and Patents Act 1988 (?)
- Users may not further distribute the material nor use it for the purposes of commercial gain.

Where a licence is displayed above, please note the terms and conditions of the licence govern your use of this document.

When citing, please reference the published version.

### **Take down policy**

While the University of Birmingham exercises care and attention in making items available there are rare occasions when an item has been uploaded in error or has been deemed to be commercially or otherwise sensitive.

If you believe that this is the case for this document, please contact [UBIRA@lists.bham.ac.uk](mailto:UBIRA@lists.bham.ac.uk) providing details and we will remove access to the work immediately and investigate.

# **Postnatal deletion of podoplanin in lymphatic endothelium results in blood filling of the lymphatic system and impairs DC migration to lymph nodes**

**Running title:** Podoplanin in adult lymphatic vessels

<sup>1</sup>Roberta Bianchi, PhD; <sup>1</sup>Erica Russo, PhD; <sup>1</sup>Samia B. Bachmann, DVM; <sup>1</sup>Steven T. Proulx, PhD; <sup>1</sup>Marko Sesartic, MSc; <sup>1</sup>Nora Smaadahl, MSc; <sup>2</sup>Steve P. Watson, PhD; <sup>3</sup>Christopher D. Buckley, MD, PhD; <sup>1</sup>Cornelia Halin, PhD and <sup>1</sup>Michael Detmar, MD\*

<sup>1</sup>Institute of Pharmaceutical Sciences, Swiss Federal Institute of Technology, ETH Zurich, Zurich, Switzerland;

<sup>2</sup>Centre for Cardiovascular Sciences, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK;

<sup>3</sup>Rheumatology Research Group, Institute for Biomedical Research, College of Medical and Dental Sciences, University of Birmingham, Birmingham, UK.

## **\*Correspondence:**

Prof. Michael Detmar  
Institute of Pharmaceutical Sciences  
Swiss Federal Institute of Technology, ETH Zurich  
Vladimir-Prelog-Weg 3, HCI H303  
8093 Zurich, Switzerland  
Tel: +41446337361  
Fax: +41446331364  
Email: [michael.detmar@pharma.ethz.ch](mailto:michael.detmar@pharma.ethz.ch)

**Keywords:** lymphatic endothelium, podoplanin, animal models

## **Subject codes:**

[129] Angiogenesis, [95] Endothelium/vascular type/nitric oxide, [97] Other Vascular biology, [98] Other Research

**Total word count:** 5624

**Total number of figures and tables:** 6

**TOC category –** Basic sciences

**TOC subcategory –** Vascular biology

## ABSTRACT

**Objective:** The lymphatic vascular system exerts major physiological functions in the transport of interstitial fluid from peripheral tissues back to the blood circulation and in the trafficking of immune cells to lymph nodes. Previous studies in global constitutive knockout mice for the lymphatic transmembrane molecule podoplanin reported perinatal lethality and a complex phenotype with lung abnormalities, cardiac defects, lymphedema, blood-filled lymphatic vessels and lack of lymph node organization, reflecting the importance of podoplanin expression not only by the lymphatic endothelium, but also by a variety of non-endothelial cell types. Therefore, we aimed to dissect the specific role of podoplanin expressed by adult lymphatic vessels.

**Approach and Results:** We generated an inducible, lymphatic-specific podoplanin knockout mouse model (Pdpn<sup>ΔLEC</sup>) and induced gene deletion postnatally. Pdpn<sup>ΔLEC</sup> mice were viable and their lymphatic vessels appeared morphologically normal with unaltered fluid drainage function. Intriguingly, Pdpn<sup>ΔLEC</sup> mice had blood-filled lymph nodes and vessels, most frequently in the neck and axillary region, and displayed a blood-filled thoracic duct, suggestive of retrograde filling of blood from the blood circulation into the lymphatic system. Histological and FACS analyses revealed normal lymph node organization with the presence of erythrocytes within lymph node lymphatic vessels but not surrounding high endothelial venules. Moreover, FITC painting experiments revealed reduced dendritic cell migration to lymph nodes in Pdpn<sup>ΔLEC</sup> mice.

**Conclusions:** These results reveal an important role of podoplanin expressed by lymphatic vessels in preventing postnatal blood filling of the lymphatic vascular system and in contributing to efficient dendritic cell migration to the lymph nodes.

## NON-STANDARD ABBREVIATIONS AND ACRONYMS

LN	lymph node
LV	lymphatic vessel
LVV	lymphovenous valve
TD	thoracic duct
LECs	lymphatic endothelial cells
FRCs	follicular reticular cells
CLEC-2	C-type lectin-like receptor 2
DCs	dendritic cells
HEVs	high endothelial venules

## INTRODUCTION

The lymphatic vascular system is responsible for tissue fluid homeostasis, immune cell trafficking and intestinal lipid uptake.<sup>1, 2</sup> Peripheral lymphatic capillaries take up interstitial fluid (i.e. lymph) that is then filtered through lymph nodes (LN) and drained by lymphatic collecting vessels that converge into the thoracic duct (TD) - the largest lymphatic vessel (LV) in the body - and the right lymphatic trunk, which finally connect to the subclavian veins, where the lymph enters the blood circulation. At these connection points, specialized valves known as lymphovenous valves (LVV) prevent the flow of blood into the lymphatic circulation.<sup>3, 4</sup> LVs also play key functions in the trafficking of immune cells from the peripheral tissues to the draining LNs, where the initiation of adaptive immune responses occurs.<sup>5</sup>

Podoplanin is a small, extensively O-glycosylated, type-1 transmembrane protein expressed by lymphatic endothelial cells (LECs)<sup>6</sup> but not by blood vascular endothelial cells and is therefore a widely used lymphatic specific marker. Podoplanin is also expressed by follicular reticular cells (FRCs) of secondary lymphoid organs<sup>7</sup>, podocytes of the kidney<sup>8</sup>, alveolar type-1 cells of the lung and by the choroid plexus epithelium<sup>9</sup>. Moreover, it is expressed in the developing heart of the mouse embryo.<sup>10</sup> Podoplanin is the only described endogenous ligand of the C-type lectin-like receptor 2 (CLEC-2)<sup>11</sup>, which is expressed by platelets<sup>12</sup> and activated dendritic cells (DCs)<sup>13-15</sup>. CLEC-2 signaling is involved in platelet activation<sup>12</sup>, DC motility<sup>15</sup> and in the maintenance of vascular integrity<sup>16</sup>, LN structure and development.<sup>17</sup>

Podoplanin global knockout mice die shortly after birth, due to a failure to inflate their lungs.<sup>18, 19</sup> Several studies described profound defects in the development and patterning of the lymphatic vasculature in these mice. Global podoplanin knockout mice display congenital lymphedema, impaired lymphatic function, dilation and mispatterning of the intestinal and cutaneous lymphatic vasculature, and blood-filled LVs.<sup>20-22</sup> The latter phenotype has been attributed to the interaction, at midgestation, of podoplanin expressed by LECs with CLEC-2 expressing platelets, inducing platelet activation and formation of thrombi that mediate the proper separation of the nascent lymphatic system from the cardinal vein.<sup>22, 23</sup> These mice also lack most of the LNs which are replaced by blood-filled, poorly organized remnants.<sup>24</sup> Additionally, several cardiac defects, including hypoplasia of the myocardium, were described in podoplanin-deficient embryos<sup>25-27</sup>, and deletion of podoplanin at the 2-cell stage causes cerebral hemorrhaging in embryos.<sup>28</sup> The broad expression of podoplanin, together with the lethal phenotype observed in global podoplanin-deficient mice, suggests an important role for podoplanin in diverse cell types and in the ontogeny of different organs. It has, however, not been possible thus far to dissect its direct function specifically in the lymphatic endothelium at late postnatal or adult stages.

To specifically investigate the function of podoplanin in the postnatal lymphatic vasculature, we developed, for the first time, an inducible, lymphatic-specific podoplanin knockout mouse by crossing *Pdpr<sup>fl/fl</sup>* mice<sup>28</sup> and *Prox1-Cre-ERT2* mice<sup>29</sup> to obtain *Pdpr<sup>ΔLEC</sup>* mice. Postnatal deletion of podoplanin was compatible with life and did not affect the maintenance of normal LV morphology and function. Strikingly, however, blood was frequently detected in the thoracic duct (TD) of *Pdpr<sup>ΔLEC</sup>* mice. Moreover, a subset of their LNs, despite maintaining a normal organization, were frequently blood-filled with detection of blood within the LVs, whereas no disruption of high endothelial venule (HEV) integrity was observed. Furthermore, DC migration to LNs after FITC challenge was reduced in *Pdpr<sup>ΔLEC</sup>* mice, highlighting a role for podoplanin in immune cell trafficking to the LN. Taken together, these results suggest important postnatal functions of LV-expressed podoplanin in preventing

retrograde flow of blood into the lymphatic system and in regulating DC migration to LNs after challenge.

## METHODS

Materials and Methods are available in the online-only Data Supplement.

## RESULTS

### *Generation of a conditional, lymphatic-specific podoplanin knockout mouse model.*

Since global constitutive podoplanin knockout mice display a lethal phenotype on most genetic backgrounds, at least in part due to a failure to inflate their lungs,<sup>18</sup> we generated mice with postnatal deletion of podoplanin specifically in the lymphatic vasculature. To this aim, Pdpn<sup>fl/fl</sup> mice<sup>28</sup> were crossed with Prox1-Cre-ERT2 mice<sup>29</sup>, which express Cre recombinase under control of the lymphatic specification gene Prox1<sup>30</sup> upon tamoxifen administration, obtaining the Pdpn<sup>ΔLEC</sup> line (Figure 1A).

Postnatal deletion of the podoplanin gene was induced by daily tamoxifen administrations from postnatal day (P) 1 to P3 (Figure 1A). Adult Pdpn<sup>ΔLEC</sup> mice showed efficient podoplanin downregulation in dermal LVs, as assessed by whole mount immunofluorescence stainings for podoplanin and the lymphatic marker LYVE-1 (Figure 1B and SID), and by FACS analysis (Figure 1C). Pdpn<sup>ΔLEC</sup> mice were monitored from birth until at least 23 weeks of age; they were not visually distinguishable from wild-type littermates and their weight was normal (data not shown).

### *Postnatal deletion of podoplanin does not interfere with lymphatic patterning.*

At P7, Pdpn<sup>ΔLEC</sup> pups appeared normal and no difference in weight was observed (Figure 2A). Podoplanin was efficiently downregulated at this time point, as assessed by podoplanin whole-mount stainings of abdominal skin, diaphragm and mesentery (Figure S1A-C). LV morphology was analyzed by staining whole-mount preparations of abdominal skin (Figure 2B), tail skin (Figure 2C) and diaphragm (Figure 2D) for LYVE-1. Morphometric analyses did not reveal any significant differences in the LYVE-1<sup>+</sup> area, number of branch points, number of ring structures, LV diameter, total vessel length and average and branch length; Figure 2 B-D).

### *Adult lymphatic vessels are morphologically and functionally normal after postnatal deletion of podoplanin.*

Next, we investigated the morphology and functionality of adult LVs lacking podoplanin. To this purpose, ear whole-mounts of 11-weeks-old mice were stained for LYVE-1. LV morphology appeared normal in adult Pdpn<sup>ΔLEC</sup> animals, as no major differences were found in the tissue area covered by LVs, the number of LV branches, the total vessel length and the average vessel diameter (Figure 3A). As a measure of lymphatic function, we injected the lymphatic specific tracer P20D800<sup>31, 32</sup> intradermally into the ear and monitored its clearance by LVs over time. As quantified by half-life and K rate of the tracer decay, no difference was observed in lymphatic clearance between Pdpn<sup>ΔLEC</sup> mice and wild-type littermates (Figure 3B).

### *Postnatal deletion of podoplanin results in blood-filled lymphatic vessels and lymph nodes.*

The LNs of adult Pdpn<sup>ΔLEC</sup> mice often displayed a bloody appearance (Figure 4A, 4C and S1A-B). In order to determine when this phenotype arose, we analyzed animals at different ages, namely P7, P21 and adults. As shown by the phenotype frequencies in different anatomical locations (Figure 4B), at P7 the LNs were never blood-filled and we only observed blood in the thoracic duct of 1 out of 9 mice analyzed. At weaning age (P21), the phenotype was apparent and, as in adult mice,

the LNs located in the neck (auricular and mandibular) and axilla (axillary and brachial) showed the phenotype with higher frequency than the inguinal LNs and flank lymphatic collectors (Figure 4C). The mesenteric LNs (Figure S1B) and intestinal LVs (Figure 4D) were blood-filled only in adult  $\text{Pdpn}^{\Delta\text{LEC}}$  mice and were not affected at P21 or P7.

FACS analysis revealed an increase in the frequency of red blood cells (identified as  $\text{CD45}^- \text{Ter119}^+$  events) in auricular LNs harvested from adult  $\text{Pdpn}^{\Delta\text{LEC}}$  mice ( $7.7 \pm 2.17\%$  of cells), as compared to wild-type littermates ( $1.3 \pm 0.53\%$ ), confirming the presence of blood in these LNs (Figure 4E).

To determine the origin of bleeding, we performed immunofluorescence analyses for red blood cells (Ter119), LYVE-1 and the pan-endothelial marker CD31 or VE-cadherin on frozen LN sections (Figure 4F and S1C-D). Red blood cells were detected inside  $\text{CD31}^+/\text{LYVE-1}^-$  HEVs (Figure 4F, arrowheads), but also inside LVs in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice (Figure 4F, arrows and high power inset, Figure S1D). Importantly, no red blood cells were observed in the proximity of HEVs, suggesting that extravasation of erythrocytes from these vessels did not occur. Since FRCs lining HEVs retained podoplanin expression in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice (Figure S1I, arrowheads), the presence of blood in the LNs was not due to compromised HEV integrity, but to blood filling of the lymphatic vasculature.

We also analyzed the presence of blood in the TD of  $\text{Pdpn}^{\Delta\text{LEC}}$  mice, which was identified by fluorescent lymphatic tracing after intradermal injection of the lymphatic tracer P20D680<sup>31</sup> in both hind paws. Blood was visible by white-light analysis in the TD in 36.4% (4 out of 11) of  $\text{Pdpn}^{\Delta\text{LEC}}$  mice (Figure 4G) but not in wild-type mice ( $n=7$ ). In the majority of  $\text{Pdpn}^{\Delta\text{LEC}}$  mice (3 out of 4), the blood-filled TD was massively dilated (Figure 4G). Importantly, only mice that showed blood in the TD also had blood-filled mesenteric LNs (Figure S1B). Taken together, these data suggest that lack of podoplanin on LECs results in retrograde flow of blood from the veins into the lymphatic system.

In order to investigate whether podoplanin is required throughout life to prevent blood-filling of the lymphatic system, we induced gene deletion in adult mice by 5 daily tamoxifen injections and analyzed the phenotype 5 days after the last tamoxifen injection (Figure S1A). With this regimen, podoplanin expression was strongly reduced even though gene deletion was not complete, as a low level of podoplanin expression was detected in some dermal LVs by whole-mount staining (Figure S1B). We did not observe any blood in the LNs, but in 1/3 mice blood filled the TD (Figure S1C). These data indicate that podoplanin is required throughout life to keep the blood from entering the lymphatic system.

*Postnatal deletion of podoplanin does not compromise lymph node organization.*

Global podoplanin knockout mice have poorly organized LNs.<sup>24</sup> To assess the contribution of LV podoplanin to LN organization, we analyzed the morphology of  $\text{Pdpn}^{\Delta\text{LEC}}$  LNs in more detail. Histological analyses revealed normal B-cell follicle organization in  $\text{Pdpn}^{\Delta\text{LEC}}$  LNs, as compared to wild-type controls (Figure 5A). Ear draining LNs of  $\text{Pdpn}^{\Delta\text{LEC}}$  mice were heavier, while the total leukocyte number measured after RBC lysis, the proportions of T cells and the absolute number of migratory DCs did not differ between the genotypes (Figure 5B-F). We also analysed HEVs by immunostaining for the differentiation marker MECA-79 and observed a normal HEV morphology in LNs of  $\text{Pdpn}^{\Delta\text{LEC}}$  mice (Figure 5G). We found heavier LNs in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice than in wild-type controls (Figure 5B). To assess the relative contribution of the presence of RBC and of the accumulation of fluid to the increased weight observed, we analysed the wet weight and dry weight of mandibular LNs and calculated the LN fluid content. Both the wet and dry LN weights

were increased in Pdpn<sup>ΔLEC</sup> mice as compared to wild-type controls, and the LN fluid content did not differ between the genotypes (Figure 5H). These data, together with the results on total LN cellularity (Figure 5C), indicate that the presence of RBC is responsible for the increased LN weight observed in Pdpn<sup>ΔLEC</sup> mice.

#### *Postnatal deletion of podoplanin impairs dendritic cell migration.*

It has been suggested that CLEC-2 deficiency hampers DC migration to the LN.<sup>15</sup> To investigate if the CLEC-2 binding partner podoplanin on LVs contributes to efficient DC trafficking to the LN, we applied FITC solution to the ears of mice and analyzed its uptake by DCs and their migration to the draining auricular LNs after 18 hours (Figure 6A). We found reduced proportions of FITC<sup>+</sup> DCs (gated as MHCII<sup>+</sup> CD11c<sup>+</sup> cells) in the LNs of Pdpn<sup>ΔLEC</sup> mice as compared to wild-type controls (Figure 6B). This reduction was a consequence of impaired DC migration, as there were no differences in the proportion of CD11b<sup>+</sup> CD11c<sup>+</sup> dermal DCs in the ear skin of Pdpn<sup>ΔLEC</sup> mice (Figure 6C). FACS analysis confirmed that LN DCs expressed CLEC-2 and that Pdpn<sup>ΔLEC</sup> mice retained normal CLEC-2 expression (Figure 6D).

## DISCUSSION

Podoplanin is a marker of lymphatic endothelium but is also expressed by other cell types such as LN FRCs and alveolar type-I cells in the lung (reviewed by Astarita et al.<sup>33</sup>). Thus, global constitutive podoplanin-deficient mice display a complex phenotype characterized by death at birth due to lung abnormalities, heart defects, lack of most LNs, edema and blood-filled LVs.<sup>18, 20, 22, 24, 26</sup> Here, for the first time, we generated and characterized a mouse model for the postnatal deletion of podoplanin specifically in LVs.

Podoplanin is the only described endogenous ligand of CLEC-2<sup>11</sup> and this biochemical interaction has been investigated in detail. The extracellular portion of podoplanin contains conserved domains required for CLEC-2 engagement<sup>34</sup>, which in platelets results in the phosphorylation of the CLEC-2 cytoplasmic tail, activation of the Syk tyrosine kinase and consequent downstream signaling via SLP-76 and PLCγ2, leading to platelet activation.<sup>11</sup> It has been proposed that during embryonic development, podoplanin/CLEC-2 induced platelet activation prevents the blood-filling of the budding lymphatic system from the cardinal vein (reviewed by Bertozzi et al.<sup>35</sup>). Indeed, mice lacking either podoplanin, CLEC-2 or downstream signaling molecules such as Syk and SLP-76, all display a similar blood-filled lymphatic phenotype.<sup>21-23, 36-39</sup> We induced podoplanin-deletion during early postnatal days (P1 to P3) since at this time point, the lymphatic system has already separated from the cardinal vein, and lymphatic plexuses are actively maturing in different organs.<sup>40</sup> We did not observe lethality or gross developmental defects up to 23 weeks of age in Pdpn<sup>ΔLEC</sup> animals, suggesting that postnatal deletion of podoplanin in lymphatic endothelium is compatible with life.

Analysis of P7 pups and adult mice revealed morphologically normal and functional LVs in Pdpn<sup>ΔLEC</sup> mice, suggesting that podoplanin does not play a major role in the postnatal maturation of LVs or in the maintenance of adult LV morphology and drainage function. Previous analyses of global constitutive podoplanin-deficient mice showed enlarged and disorganized LVs, as well as edema and impaired lymphatic function.<sup>20</sup> Additionally, blocking of podoplanin function by administration of podoplanin-Fc during late embryonic development resulted in a less complex diaphragmatic lymphatic network in pups.<sup>41</sup> Thus, it is likely that, early in development, podoplanin functionally contributes not only to the separation of the blood vascular and the lymphatic system, but also to LV patterning. However, our data indicate that after birth, podoplanin is dispensable for proper lymphatic vessel maintenance and drainage function. In global podoplanin-deficient mice, where

podoplanin is also deleted from LN FRCs, lymphatic function is likely impaired by the lack of LNs.<sup>24</sup> In contrast, in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice, LNs are present and are normally organized, indicating that podoplanin expressed by FRCs and not by LECs is responsible for the lack of LN organization observed in global constitutive podoplanin-deficient mice.

The most striking phenotype observed in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice was the presence of blood-filled LN LVs especially in the neck and axilla (95.5% and 68.2% of adult  $\text{Pdpn}^{\Delta\text{LEC}}$  mice respectively) and, less frequently, flank collecting LVs (7.6% of adult  $\text{Pdpn}^{\Delta\text{LEC}}$  mice) and mesenteric LNs and intestinal LVs (31.8% and 10.6% of adult  $\text{Pdpn}^{\Delta\text{LEC}}$  mice respectively). Moreover, blood was present in the TD of 36.4% of adult  $\text{Pdpn}^{\Delta\text{LEC}}$  mice analyzed. Interestingly, LN LVs were never blood-filled in P7  $\text{Pdpn}^{\Delta\text{LEC}}$  mice (i.e., 5 days after the last tamoxifen injection) and only 11% of mice showed a blood-filled TD at this time point. At weaning age (P21), the frequency of the phenotype was similar to adult mice, with the exception of the mesenteric LNs and LVs, which were never blood-filled at this time point. These data suggest that, after podoplanin deletion, the blood takes a few weeks to reach the LN LVs and more distal sites, such as the mesentery. Accordingly, when gene deletion was induced in adult mice, LN LVs were not blood-filled 5 days after the last tamoxifen injection, but blood was detected in the TD of 1/3 of the mice, similarly to what we observed at P7 after tamoxifen injections at P1 to P3.

Moreover, the incomplete penetrance of this phenotype might be due to partial podoplanin deletion, which can occur when Cre recombination is induced after embryonic development.<sup>42</sup> Efficient gene deletion relies on sufficient tamoxifen bioavailability in a given cell/organ, Cre promoter activity and expression levels of the target gene and local chromatin conformation at the loxP sites, and it is possible that these parameters vary between different cells.<sup>42-44</sup>

Taken together, these data allowed us to propose a potential mechanism of the blood filling of LVs in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice. The observation that the LNs located in the neck and axilla, which are anatomically closer to the LVVs, were filled by blood with the highest frequency, and that the phenotype was more pronounced in adult mice than at P7, suggests that LV expression of podoplanin regulates blood-lymph separation in adults and that loss of podoplanin likely compromises platelet aggregation in the proximity of the LVV, allowing filling of blood into the lymphatic system (Figure SV). Our observations confirm and extend a recent study that revealed an important function of platelet CLEC-2 in lymphovenous hemostasis.<sup>45</sup> Loss of CLEC-2 in the hematopoietic lineage using fetal liver chimeras, genetic approaches or antibody treatment caused filling of the TD with blood from the subclavian vein; thus, these mice displayed blood-filled mesenteric LNs and vessels.<sup>45, 46</sup> Our study, for the first time, directly addresses the important role that podoplanin on LECs, as the molecular counterpart of CLEC-2, plays in controlling blood-lymphatic separation in developed animals.

Several reports described blood-filled LNs and Peyer's patches in adult mice lacking CLEC-2 specifically in the megakaryocyte/platelet lineage or after treatment with CLEC-2 depleting antibodies.<sup>16, 17, 45</sup> However, in models of CLEC-2 deficiency, LN bleeding was also described to occur due to compromised HEV integrity.<sup>16</sup> It was suggested that podoplanin on FRCs interacts with CLEC-2 expressing platelets, inducing release of sphingosine-1-phosphate that tightens inter-endothelial junctions in HEVs.<sup>16</sup> Our immunofluorescence analyses of frozen LN sections in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice revealed blood cells inside LVs of the LNs, but never in close proximity to HEVs. Moreover, FRCs normally lined HEVs and retained podoplanin expression in  $\text{Pdpn}^{\Delta\text{LEC}}$  LNs. These observations indicate that the blood-filled LN phenotype was



not caused by disruption of HEV integrity, but solely by blood filling into the LVs of the LNs.

The direct analysis of the LVV and the presence of thrombi in its proximity has been recently studied by whole-embryo histology<sup>45</sup>. However, the investigation of this anatomical structure by histology in adult animals is technically challenging due to its deep location in the upper chest, and to the best of our knowledge, has not been described so far. Our detailed anatomical analysis of the affected LNs and major lymphatic vessels (i.e. the TD) that constitute the terminal end of the lymphatic system, together with the published data of models of CLEC-2 deficiency<sup>45</sup>, strongly support our hypothesis that podoplanin functions at the interface between the lymphatic and the blood vascular system to prevent flow of blood into the lymphatic circulation postnatally (Figure SV).

Despite the presence of blood-filled LVs in the LN, the lymphatic drainage function was not compromised in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice under steady-state conditions, as evaluated by the clearance of an injected tracer from ear skin. Moreover, we could perfuse a lymphatic tracer into the blood-filled TD of  $\text{Pdpn}^{\Delta\text{LEC}}$  mice after intradermal injections into the hind paws. Similarly, Hess and colleagues observed fluorescence in the TD of CLEC-2 deficient animals after FITC-dextran injection in the hind limb, suggesting that forward lymphatic flow was present even when blood filled the lymphatic system in a model of CLEC-2 deficiency.<sup>45</sup> It is possible that this residual forward flow is sufficient to ensure physiological fluid drainage in  $\text{Pdpn}^{\Delta\text{LEC}}$  and it would be of interest to test the drainage function in conditions characterized by increased demand of fluid drainage, such as inflammation. Another possible explanation could be that new alternate flow routes may be used to bypass the affected LNs, similarly to what has been described in metastasis bearing LNs.<sup>31</sup>

Surprisingly,  $\text{Pdpn}^{\Delta\text{LEC}}$  mice do not show signs of distress or lethality, despite the presence of blood in the LVs. It is not uncommon that the presence of profound lymphatic defects does not translate in lethality; a remarkable example is the K14-sVEGFR3 mouse line, which is viable despite completely lacking dermal LVs and displaying lymphedema<sup>47</sup>.

Importantly, we found a reduced DC migration to LNs in response to FITC challenge in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice. It has been reported that CLEC-2 expressed by DCs interacts with podoplanin expressed by LECs and FRCs to induce actin cytoskeleton rearrangement and cell motility.<sup>15</sup> Consequently, CLEC-2<sup>-/-</sup> fetal liver chimeras or mice lacking CLEC-2 specifically in CD11c<sup>+</sup> DCs show reduced proportions of DCs migrating to the LNs.<sup>15, 48</sup> Here, we complement these observations made in CLEC-2 deficient mice and reinforce the importance of podoplanin expressed by LVs in DC migration. Importantly, we observed no differences in the number of migratory DCs present in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice under steady-state conditions. Similarly, HEV maintenance, which has been linked to DC migration<sup>49</sup>, was not affected. These data suggest that CLEC-2 impacts on DC migration upon induction of inflammation and they are in agreement with published data showing no constitutive CLEC-2 expression in resting DCs, but upregulation upon LPS-induced systemic inflammation in a small subset of activated DCs.<sup>14</sup>

Taken together, our study highlights an important role of podoplanin in adult lymphatic vessels: podoplanin prevents blood mixing into the lymphatic system postnatally and contributes to efficient DC migration to the lymph node in response to inflammation.

## ACKNOWLEDGEMENTS

We thank Taija Mäkinen (Uppsala University) for kindly providing the Prox1-Cre-ERT2 mouse line; Carlos Ochoa, Sven Novok and Marion Hermerschmidt for animal caretaking; Sinem Karaman and Alexandra M. Ochsenbein for helpful discussions and valuable help; Jeannette Scholl, Peter Camenzind and Yi-Fan Lim for excellent technical assistance and the Scientific Center for Optical and Electron Microscopy (ScopeM) of ETH Zurich for support with confocal microscopy.

## SOURCE OF FUNDING

This work was supported by the Swiss National Science Foundation Grant 31003A-130627, European Research Council Grant LYVICAM, Oncosuisse, Krebsliga Zurich, the Leducq Foundation (to MD), and by the British Heart Foundation CH/03/003 (to SPW).

## DISCLOSURES

None.

## REFERENCES

1. Cueni LN, Detmar M. The lymphatic system in health and disease. *Lymphat Res Biol.* 2008;6:109-122
2. Alitalo K. The lymphatic vasculature in disease. *Nat Med.* 2011;17:1371-1380
3. El Zawahry MD, Sayed NM, El-Awady HM, Abdel-Latif A, El-Gindy M. A study of the gross, microscopic and functional anatomy of the thoracic duct and the lympho-venous junction. *Int Surg.* 1983;68:135-138
4. Srinivasan RS, Oliver G. Prox1 dosage controls the number of lymphatic endothelial cell progenitors and the formation of the lymphovenous valves. *Genes Dev.* 2011;25:2187-2197
5. Randolph GJ, Angeli V, Swartz MA. Dendritic-cell trafficking to lymph nodes through lymphatic vessels. *Nat Rev Immunol.* 2005;5:617-628
6. Breiteneder-Geleff S, Soleiman A, Kowalski H, Horvat R, Amann G, Kriehuber E, Diem K, Weninger W, Tschachler E, Alitalo K, Kerjaschki D. Angiosarcomas express mixed endothelial phenotypes of blood and lymphatic capillaries: Podoplanin as a specific marker for lymphatic endothelium. *Am J Pathol.* 1999;154:385-394
7. Farr AG, Berry ML, Kim A, Nelson AJ, Welch MP, Aruffo A. Characterization and cloning of a novel glycoprotein expressed by stromal cells in t-dependent areas of peripheral lymphoid tissues. *J Exp Med.* 1992;176:1477-1482
8. Breiteneder-Geleff S, Matsui K, Soleiman A, Meraner P, Poczewski H, Kalt R, Schaffner G, Kerjaschki D. Podoplanin, novel 43-kd membrane protein of glomerular epithelial cells, is down-regulated in puromycin nephrosis. *Am J Pathol.* 1997;151:1141-1152
9. Williams MC, Cao Y, Hinds A, Rishi AK, Wetterwald A. T1 alpha protein is developmentally regulated and expressed by alveolar type i cells, choroid plexus, and ciliary epithelia of adult rats. *Am J Respir Cell Mol Biol.* 1996;14:577-585
10. Gittenberger-de Groot AC, Mahtab EA, Hahurij ND, Wisse LJ, Deruiter MC, Wijffels MC, Poelmann RE. Nkx2.5-negative myocardium of the posterior heart field and its correlation with podoplanin expression in cells from the developing cardiac pacemaking and conduction system. *Anat Rec.* 2007;290:115-122
11. Suzuki-Inoue K, Kato Y, Inoue O, Kaneko MK, Mishima K, Yatomi Y, Yamazaki Y, Narimatsu H, Ozaki Y. Involvement of the snake toxin receptor clec-2, in podoplanin-mediated platelet activation, by cancer cells. *J Biol Chem.* 2007;282:25993-26001

12. Suzuki-Inoue K, Fuller GL, Garcia A, et al. A novel syk-dependent mechanism of platelet activation by the c-type lectin receptor clec-2. *Blood*. 2006;107:542-549
13. Colonna M, Samaridis J, Angman L. Molecular characterization of two novel c-type lectin-like receptors, one of which is selectively expressed in human dendritic cells. *Eur J Immunol*. 2000;30:697-704
14. Lowe KL, Navarro-Nunez L, Benezech C, Nayar S, Kingston BL, Nieswandt B, Barone F, Watson SP, Buckley CD, Desanti GE. The expression of mouse clec-2 on leucocyte subsets varies according to their anatomical location and inflammatory state. *Eur J Immunol*. 2015;45:2484-2493
15. Acton S, Astarita JL, Malhotra D, et al. Podoplanin-rich stromal networks induce dendritic cell motility via activation of the c-type lectin receptor clec-2. *Immunity*. 2012;37:1-14
16. Herzog BH, Fu J, Wilson SJ, et al. Podoplanin maintains high endothelial venule integrity by interacting with platelet clec-2. *Nature*. 2013;502:105-109
17. Benezech C, Nayar S, Finney BA, Withers DR, Lowe K, Desanti GE, Marriott CL, Watson SP, Caamano JH, Buckley CD, Barone F. Clec-2 is required for development and maintenance of lymph nodes. *Blood*. 2014;123:3200-3207
18. Ramirez MI, Millien G, Hinds A, Cao Y, Seldin DC, Williams MC. T1alpha, a lung type i cell differentiation gene, is required for normal lung cell proliferation and alveolus formation at birth. *Dev Biol*. 2003;256:61-72
19. Millien G, Spira A, Hinds A, Wang J, Williams MC, Ramirez MI. Alterations in gene expression in t1 alpha null lung: A model of deficient alveolar sac development. *BMC Dev Biol*. 2006;6:35
20. Schacht V, Ramirez MI, Hong Y-K, Hirakawa S, Feng D, Harvey N, Williams M, Dvorak AM, Dvorak HF, Oliver G, Detmar M. T1alpha/podoplanin deficiency disrupts normal lymphatic vasculature formation and causes lymphedema. *EMBO J*. 2003;22:3546-3556
21. Fu J, Gerhardt H, McDaniel JM, et al. Endothelial cell o-glycan deficiency causes blood/lymphatic misconnections and consequent fatty liver disease in mice. *J Clin Invest*. 2008;118:3725-3737
22. Uhrin P, Zaujec J, Breuss JM, Olcaydu D, Chrenek P, Stockinger H, Fuerthbauer E, Moser M, Haiko P, Fässler R, Alitalo K, Binder BR, Kerjaschki D. Novel function for blood platelets and podoplanin in developmental separation of blood and lymphatic circulation. *Blood*. 2010;115:3997-4005
23. Bertozzi CC, Schmaier AA, Mericko P, et al. Platelets regulate lymphatic vascular development through clec-2-slp-76 signaling. *Blood*. 2010;116:661-670
24. Peters A, Pitcher LA, Sullivan JM, Mitsdoerffer M, Acton SE, Franz B, Wucherpfennig K, Turley S, Carroll MC, Sobel RA, Bettelli E, Kuchroo VK. Th17 cells induce ectopic lymphoid follicles in central nervous system tissue inflammation. *Immunity*. 2011;35:986-996
25. Mahtab EA, Vicente-Steijn R, Hahurij ND, Jongbloed MR, Wisse LJ, DeRuiter MC, Uhrin P, Zaujec J, Binder BR, Schalijs MJ, Poelmann RE, Gittenberger-de Groot AC. Podoplanin deficient mice show a rhoa-related hypoplasia of the sinus venosus myocardium including the sinoatrial node. *Dev Dyn*. 2009;238:183-193
26. Mahtab EA, Wijffels MC, Van Den Akker NM, Hahurij ND, Lie-Venema H, Wisse LJ, Deruiter MC, Uhrin P, Zaujec J, Binder BR, Schalijs MJ, Poelmann RE, Gittenberger-De Groot AC. Cardiac malformations and myocardial abnormalities in podoplanin knockout mouse embryos: Correlation with abnormal epicardial development. *Dev Dyn*. 2008;237:847-857
27. Douglas YL, Mahtab EA, Jongbloed MR, Uhrin P, Zaujec J, Binder BR, Schalijs MJ, Poelmann RE, Deruiter MC, Gittenberger-de Groot AC. Pulmonary vein, dorsal atrial wall and atrial septum abnormalities in

- podoplanin knockout mice with disturbed posterior heart field contribution. *Pediatr Res*. 2009;65:27-32
28. Lowe KL, Finney BA, Deppermann C, Hagerling R, Gazit SL, Frampton J, Buckley C, Camerer E, Nieswandt B, Kiefer F, Watson SP. Podoplanin and clec-2 drive cerebrovascular patterning and integrity during development. *Blood*. 2015;125:3769-3777
  29. Bazigou E, Lyons OT, Smith A, Venn GE, Cope C, Brown NA, Makinen T. Genes regulating lymphangiogenesis control venous valve formation and maintenance in mice. *J Clin Invest*. 2011;121:2984-2992
  30. Wigle JT, Oliver G. Prox1 function is required for the development of the murine lymphatic system. *Cell*. 1999;98:769-778
  31. Proulx ST, Luciani P, Christiansen A, Karaman S, Blum KS, Rinderknecht M, Leroux JC, Detmar M. Use of a peg-conjugated bright near-infrared dye for functional imaging of rerouting of tumor lymphatic drainage after sentinel lymph node metastasis. *Biomaterials*. 2013;34:5128-5137
  32. Karaman S, Buschle D, Luciani P, Leroux JC, Detmar M, Proulx ST. Decline of lymphatic vessel density and function in murine skin during aging. *Angiogenesis*. 2015;18:489-498
  33. Astarita JL, Acton SE, Turley SJ. Podoplanin: Emerging functions in development, the immune system, and cancer. *Front Immunol*. 2012;3:283
  34. Kato Y, Fujita N, Kunita A, Sato S, Kaneko M, Osawa M, Tsuruo T. Molecular identification of aggrus/t1alpha as a platelet aggregation-inducing factor expressed in colorectal tumors. *J Biol Chem*. 2003;278:51599-51605
  35. Bertozzi CC, Hess PR, Kahn ML. Platelets: Covert regulators of lymphatic development. *Arterioscler Thromb Vasc Biol*. 2010;30:2368-2371
  36. Abtahian F, Guerriero A, Sebzda E, et al. Regulation of blood and lymphatic vascular separation by signaling proteins slp-76 and syk. *Science*. 2003;299:247-251
  37. Carramolino L, Fuentes J, Garcia-Andres C, Azcoitia V, Riethmacher D, Torres M. Platelets play an essential role in separating the blood and lymphatic vasculatures during embryonic angiogenesis. *Circ Res*. 2010;106:1197-1201
  38. Finney BA, Schweighoffer E, Navarro-Núñez L, et al. Clec-2 and syk in the megakaryocytic/platelet lineage are essential for development. *Blood*. 2012;119:1747-1756
  39. Osada M, Inoue O, Ding G, Shirai T, Ichise H, Hirayama K, Takano K, Yatomi Y, Hirashima M, Fujii H, Suzuki-Inoue K, Ozaki Y. Platelet activation receptor clec-2 regulates blood/lymphatic vessel separation by inhibiting proliferation, migration, and tube formation of lymphatic endothelial cells. *J Biol Chem*. 2012;287:22241-22252
  40. Yang Y, Oliver G. Development of the mammalian lymphatic vasculature. *J Clin Invest*. 2014;124:888-897
  41. Bianchi R, Fischer E, Yuen D, Ernst E, Chen L, Otto VI, Detmar M. Mutation of threonin 34 in mouse podoplanin-fc reduces clec-2 binding and toxicity in vivo while retaining anti-lymphangiogenic activity. *J Biol Chem*. 2014;289:21016-21027
  42. Feil S, Valtcheva N, Feil R. Inducible cre mice. *Methods Mol Biol*. 2009;530:343-363
  43. Monvoisin A, Alva JA, Hofmann JJ, Zovein AC, Lane TF, Iruela-Arispe ML. Ve-cadherin-creert2 transgenic mouse: A model for inducible recombination in the endothelium. *Dev Dyn*. 2006;235:3413-3422
  44. Vooijs M, Jonkers J, Berns A. A highly efficient ligand-regulated cre recombinase mouse line shows that loxp recombination is position dependent. *EMBO Rep*. 2001;2:292-297

45. Hess PR, Rawnsley DR, Jakus Z, Yang Y, Sweet DT, Fu J, Herzog B, Lu M, Nieswandt B, Oliver G, Makinen T, Xia L, Kahn ML. Platelets mediate lymphovenous hemostasis to maintain blood-lymphatic separation throughout life. *J Clin Invest.* 2014;124:273-284
46. Boulaftali Y, Hess PR, Kahn ML, Bergmeier W. Platelet immunoreceptor tyrosine-based activation motif (itam) signaling and vascular integrity. *Circ Res.* 2014;114:1174-1184
47. Makinen T, Jussila L, Veikkola T, Karpanen T, Kettunen MI, Pulkkanen KJ, Kauppinen R, Jackson DG, Kubo H, Nishikawa S, Yla-Herttuala S, Alitalo K. Inhibition of lymphangiogenesis with resulting lymphedema in transgenic mice expressing soluble vegf receptor-3. *Nat Med.* 2001;7:199-205
48. Astarita JL, Cremasco V, Fu J, et al. The clec-2-podoplanin axis controls the contractility of fibroblastic reticular cells and lymph node microarchitecture. *Nat Immunol.* 2015;16:75-84
49. Moussion C, Girard JP. Dendritic cells control lymphocyte entry to lymph nodes through high endothelial venules. *Nature.* 2011;479:542-546

## HIGHLIGHTS

- Generation of the first inducible, lymphatic-specific, podoplanin knockout mouse model (Pdpn<sup>ΔLEC</sup> mice)
- Postnatal loss of podoplanin in lymphatic endothelium results in blood-filled lymphatic vessels and lymph nodes without affecting lymphatic vessel patterning, morphology and drainage function of lymphatic vessels or lymph node organization
- Pdpn<sup>ΔLEC</sup> mice show impaired dendritic cell migration to lymph nodes
- Podoplanin on adult lymphatic vessels plays important functions in maintaining lymph-blood separation and in contributing to efficient dendritic cell migration.

## FIGURE LEGENDS

**Figure 1. Generation of conditional, lymphatic-specific podoplanin knockout mice.** **A.** Pdpn<sup>fl/fl</sup> mice were crossed with Prox1-Cre-ERT2 mice to obtain the Pdpn<sup>ΔLEC</sup> line. Postnatal gene deletion was induced by tamoxifen administration from postnatal day (P)1 to P3. **B.** Confocal z-stack images of whole-mount ear skin samples show that podoplanin was undetectable on LYVE-1<sup>+</sup> LVs in adult Pdpn<sup>ΔLEC</sup> animals. Scale bars: 100 μm. **C.** Ear skin single cell suspensions were analyzed by FACS. LECs were gated as CD45<sup>-</sup> CD31<sup>+</sup> LYVE-1<sup>+</sup> events. Representative histograms of podoplanin expression on LECs isolated from adult Pdpn<sup>ΔLEC</sup> and wild-type control mice are shown. The median fluorescence intensity (MFI) for podoplanin in the LEC gate was significantly lower in Pdpn<sup>ΔLEC</sup> samples. Mann-Whitney test, *P*<0.001. At least 6 mice per genotype were analyzed.

**Figure 2. Postnatal deletion of podoplanin does not interfere with lymphatic patterning.** **A.** Comparable body weights of Pdpn<sup>ΔLEC</sup> mice and wild-type littermates at P7. Unpaired Student's t-test, *P*>0.05. **B.** Confocal z-stack images of representative whole-mounts of abdominal skin samples stained for LYVE-1 (green). Quantification of the LYVE-1<sup>+</sup> area and number of branch points per image showed no difference between genotypes. Mann-Whitney test, *P*>0.05. **C.** Confocal z-stack images of representative tail skin whole-mount samples stained for LYVE-1. Quantification of the LYVE-1<sup>+</sup> area and the number of ring-like structures per image revealed no difference between genotypes. Mann-Whitney test, *P*>0.05. **D.** Confocal z-stack images of representative diaphragm segments stained for LYVE-1. Quantification of the LYVE-1<sup>+</sup> area, number of branch points, total vessel length,

average vessel diameter and average branch length per diaphragmatic segment showed no difference between genotypes. Mann-Whitney test,  $P>0.05$ . Four to five animals per genotype from 4 independent litters were analyzed. Scale bars: 100  $\mu\text{m}$ .

**Figure 3. Adult lymphatic vessels are morphologically and functionally normal after postnatal deletion of podoplanin.** **A.** Confocal z-stack images of ear skin samples whole-mounts stained for LYVE-1 (red). Scale bar: 100  $\mu\text{m}$ . Quantification of the LYVE-1<sup>+</sup> area, number of branches, total vessel length and average vessel diameter showed no difference between genotypes. Mann-Whitney test,  $P>0.05$ . Six animals per genotype from independent litters were analyzed. **B.** IVIS images of ears injected intradermally with P20D800 tracer at several time points after injection. Clearance of the tracer by LVs was calculated as half-life and K rate and showed no difference between the two groups. Unpaired Student's t-test,  $P>0.05$ . Each data point represents one injected ear. Six to eight animals per genotype were analyzed.

**Figure 4. Postnatal deletion of podoplanin results in blood-filled LVs and lymph nodes.** **A.** Photograph of auricular LNs harvested from adult mice.  $\text{Pdpn}^{\Delta\text{LEC}}$  LNs appear blood-filled. Scale bar: 1 mm. **B.** Frequency of blood-filled LNs of the neck (auricular and mandibular), axilla (axillary and brachial), mesentery (mes) and inguinal (ing) regions and thoracic duct (TD) in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice at different time points (P7 n=9; P21 n=5; adults n=66, n=11 for TD analysis; nd: not determined). **C.** Normal anatomy of the lymphatic flank collector and representative photograph of a  $\text{Pdpn}^{\Delta\text{LEC}}$  mouse displaying filling of the collector (arrowheads) with blood, from the axillary (ax) LN to the inguinal (ing) LN. Scale bars: 5 mm. **D.** Normal anatomy of the adult mesentery and representative photograph of a  $\text{Pdpn}^{\Delta\text{LEC}}$  mouse with blood-filled intestinal LVs. Scale bars: 1 mm (inset: 100  $\mu\text{m}$ ). **E.** Representative FACS plots of single cell LN suspensions stained for the red blood cell marker Ter119 and the leukocyte marker CD45. The frequency of red blood cells in  $\text{Pdpn}^{\Delta\text{LEC}}$  LNs was significantly higher than in wild-type LNs. Mann-Whitney test,  $**P<0.01$ . **F.** Confocal images of frozen sections of axillary LN stained for LYVE-1 (green), Ter119 (red) and CD31 (cyan). Red blood cells (red) filled LN LVs (green; arrows) in  $\text{Pdpn}^{\Delta\text{LEC}}$  samples. Red blood cells were also found inside HEVs (cyan; arrowheads) but never in their proximity. A minimum of 6 animals per genotype and condition were analyzed. Boxed area corresponds to higher power inset. Scale bars: 100  $\mu\text{m}$  (inset: 25  $\mu\text{m}$ ). **G.** The fluorescent tracer P20D680 was used to identify the thoracic duct after injection into both hind paws. Blood was present in the TD of 36.4 % of  $\text{Pdpn}^{\Delta\text{LEC}}$  mice, scale bar: 1 mm.

**Figure 5. Postnatal deletion of podoplanin does not compromise lymph node organization.** **A.** Confocal tile-scan of axillary LNs stained for the B cell marker B220 and with DAPI shows normal B-cell follicle organization in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice. Scale bars: 200  $\mu\text{m}$ . **B.** Auricular LNs were significantly heavier in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice. Unpaired Student's t-test,  $**P<0.01$ . **C.** Total cellularity after red blood cell lysis showed no difference between genotypes. Mann-Whitney test,  $P>0.05$ . **D.** Total number of leukocytes (CD45<sup>+</sup> cells) per LN was not different between genotypes. Mann-Whitney test,  $P>0.05$ . **E.** Proportions of T cells (% of CD3<sup>+</sup> cells in CD45<sup>+</sup> gate) did not differ between genotypes. Mann-Whitney test,  $P>0.05$ . **F.** Total number of migratory DCs (CD45<sup>+</sup> MHCII<sup>high</sup> CD11c<sup>+</sup> cells) did not differ between genotypes. Mann-Whitney test,  $P>0.05$ . **G.** Representative confocal images of LN frozen sections stained for MECA79 (red) and podoplanin (green) revealed normal HEVs in  $\text{Pdpn}^{\Delta\text{LEC}}$  mice. Scale bars: 10  $\mu\text{m}$ . **H.** Wet and dry weight of mandibular LNs were measured to calculate the fluid content. There was no difference in LN fluid content between the genotypes. Unpaired Student's t-test,  $P>0.05$ .

**Figure 6. Postnatal deletion of podoplanin impairs dendritic cell migration to LNs.** **A.** Schematic representation of the FITC painting experiment: FITC solution was applied to the ears, and the draining auricular LNs were harvested and analyzed by FACS after 18 hours. **B.** Representative FACS dot plots (gated on MHCII<sup>+</sup> events) showing MHCII<sup>+</sup> CD11c<sup>+</sup> FITC<sup>+</sup> DCs in wild-type and Pdpn<sup>ΔLEC</sup> mice 18 hours after FITC application. Quantification of the proportions of MHCII<sup>+</sup> CD11c<sup>+</sup> FITC<sup>+</sup> DCs in the auricular draining LNs 18 hours after FITC application showed a significant reduction in Pdpn<sup>ΔLEC</sup> mice. Mann-Whitney test, \**P*<0.05. Pooled data from 2 independent experiments analyzing a total of 7 mice per genotype are shown. **C.** The proportions of CD45<sup>+</sup> CD11b<sup>+</sup> CD11c<sup>+</sup> DCs in the ear skin did not differ between Pdpn<sup>ΔLEC</sup> and wild-type control mice. Mann-Whitney test, *P*>0.05. Left panel shows pooled data from 3 independent experiments analyzing a total of 11 mice per genotype. Right panel shows the mean value obtained in the 3 experiments. **D.** FACS analysis shows surface CLEC-2 expression on LN DCs (gated on CD45<sup>+</sup> MHCII<sup>+</sup> CD11c<sup>+</sup> cells), histograms and delta median fluorescent intensities are shown. Wild-type and Pdpn<sup>ΔLEC</sup> mice express comparable levels of CLEC-2 (n=5).